

Product Information	
End Repair Mix	
Part Number	Y9140-LC-L
Concentration	1 Rxn/ μ L
Unit Size	200 Reactions
Storage Temperature	-25°C to -15°C
Lot Number	
Reference Number	

Product Description: The End-Repair mix converts DNA containing damaged or incompatible 5'- and/or 3'- protruding ends to 5'-phosphorylated, blunt-ended DNA. This low-concentration formulation of the End-Repair Mix is compatible with applications requiring <1 microgram of DNA to be prepared for blunt-end ligation. The conversion to blunt-ended DNA is accomplished by exploiting the 5'→3' polymerase and 3'→5' exonuclease activities of T4 DNA Polymerase (P7080). T4 Polynucleotide Kinase (Y9040) ensures that the ends of the blunt-ended DNA fragments are 5'-phosphorylated for subsequent ligation by T4 DNA Ligase (L6030-HC).

Product Specifications						
Y9140						
Assay	SDS Purity	3'→5' Nuclease	5' Phosphorylation	5'→3' DNA Synthesis	DS Endonuclease	E. coli DNA Contamination
Units Tested	n/a	n/a	n/a	n/a	10 μ L	10 μ L
Specification	>99%	Functional	Functional	Functional	No Conversion	<10 copies

Source of Protein: Purified from strains of E. coli that express the recombinant T4 DNA Polymerase, and T4 Polynucleotide Kinase genes, respectively.

Quality Control Analysis:

Functionality is assessed by adding 2 μ L of End-Repair Mix to a double restriction enzyme digested, dephosphorylated plasmid DNA in 1X reaction buffer containing 0.1mM dNTPs and incubated at 25°C for 30 minutes. Competent cells were transformed with the ligation mixture, plated onto LB/Amp/X-Gal plates and incubated overnight at 37°C. Control reactions consisting of End-Repair Mix without T4 DNA polymerase and/or T4 Polynucleotide Kinase were tested in parallel. The efficiency of the reaction was evaluated by comparing the number of blue and white colonies present in the End-Repair Mix plates to those of the control plates.

Contamination Tests:

Purified free of contaminating endonucleases. In addition, >99% enzyme purity is analyzed by SDS-PAGE, and negligible E.coli genomic DNA is confirmed by qPCR.

Supplied in: 10mM Tris-HCl, 100 mM KCl, 1mM DTT, 0.1mM EDTA, < 0.1% Triton X-100, 50% glycerol pH 7.4 @ 25°C.

Supplied with:

10X End-Repair Buffer (B9140): 1M Tris-HCl, 500mM NaCl, 100mM MgCl₂, 50mM DTT, 0.25% Triton X-100, pH 7.5 @ 25°C.
N2060 (1mM dNTPs)

Notes:

ATP is not required because the T4 Polynucleotide Kinase can utilize the deoxynucleotides (dATP and dTTP) used in the reaction as phosphate donors.

Limitations of Use

This product was developed, manufactured, and sold for *in vitro* use only. The product is not suitable for administration to humans or animals. SDS sheets relevant to this product are available upon request.

Usage Instructions:

1. Purify DNA to be blunted, dissolve in TE buffer.
2. Combine and mix the following components in a sterile tube:
 - 1-19 μL Purified DNA (up to 5 μg)
 - 2.5 μL 10X End-Repair Buffer
 - 2.5 μL 1 mM dNTP mix (N2060)
 - 1-3 μL End-Repair Enzyme Mix
 - Sterile H_2O to 25 μL
 - Total Volume: 25 μL**
3. Incubate room temperature (25°C) 30 minutes.
Inactivate End-Repair Enzyme by heat at 75°C for 20 minutes.
4. Ligation may be performed immediately using Enzymatics Rapid format T4 DNA Ligase (L6030-HC).

Limitations of Use

This product was developed, manufactured, and sold for *in vitro* use only. The product is not suitable for administration to humans or animals. SDS sheets relevant to this product are available upon request.